

ITP-SB-93-37
hep-ph/9307344
July, 1993

A New Model for Fermion Masses in Supersymmetric Grand Unified Theories

Alexander Kusenko¹ and Robert Shrock²

Institute for Theoretical Physics
State University of New York
Stony Brook, N. Y. 11794-3840

Abstract

We present a simple model for fermion mass matrices and quark mixing in the context of supersymmetric grand unified theories and show its agreement with experiment. Our model realizes the GUT mass relations $m_d = 3m_e$, $m_s = m_\mu/3$, $m_b = m_\tau$ in a new way and is easily consistent with values of m_t suggested by MSSM fits to LEP data.

¹email: sasha@max.physics.sunysb.edu

²email: shrock@max.physics.sunysb.edu; partially supported by NSF grant PHY-91-08054

Understanding the values of fermion masses and quark mixing remains one of the most important unsolved problems in particle physics. The magnitude of the challenge is shown by the fact that at present it is not definitely clear which approach one should take to the related matter of electroweak symmetry breaking (EWSB). Intensive efforts continue in at least two quite different directions: (i) supersymmetry, and, in particular, the minimal supersymmetric standard model (MSSM), in which EWSB is perturbative [1]; (ii) dynamical (and nonperturbative) EWSB, e.g. slowly running technicolor [2], as well as others such as compositeness. Here we shall adopt the first approach, specifically the MSSM, which may provide the simplest way to understand the precision electroweak measurements from LEP [3] while stabilizing the Higgs sector in the standard model. A further motivation for the MSSM is that it achieves unification of gauge couplings at a single scale m_G , a prerequisite for a supersymmetric grand unified theory (SGUT).

In GUT's such as SU(5) or SO(10), by well-known restrictions on (renormalizable) Higgs couplings, the elements of the lepton Yukawa matrix can be made equal to 1 or -3 times corresponding elements of down quark Yukawa matrix. In this approach, one can get a simple relation between the resultant (running) lepton and down quark masses, evaluated at the GUT scale. One interesting proposal [4] is that at the GUT scale

$$m_d = 3m_e, \quad m_s = \frac{m_\mu}{3}, \quad m_b = m_\tau \quad (1)$$

It is important to determine which (experimentally acceptable) forms for fermion mass matrices yield this relation. Since 1979, essentially only one such form (up to trivial equivalence transformations) has been found. With $-\mathcal{L}_m = v_u \bar{\psi}_{u,L} Y_u \psi_{u,R} + v_d (\bar{\psi}_{d,L} Y_d \psi_{d,R} + \bar{\psi}_{e,L} Y_e \psi_{e,R}) + h.c.$, this form has GUT scale Yukawa matrices

$$Y_u = \begin{pmatrix} 0 & A_u & 0 \\ A_u & 0 & B_u \\ 0 & B_u & C_u \end{pmatrix} \quad (2)$$

$$Y_d = \begin{pmatrix} 0 & A_d e^{i\phi} & 0 \\ A_d e^{-i\phi} & B_d & 0 \\ 0 & 0 & C_d \end{pmatrix} \quad (3)$$

$$Y_e = \begin{pmatrix} 0 & A_d & 0 \\ A_d & -3B_d & 0 \\ 0 & 0 & C_d \end{pmatrix} \quad (4)$$

Interesting studies of SGUT Yukawa matrices, and this ansatz in particular, have been carried out recently using renormalization group equations (RGE's) in the MSSM [5, 6, 7] (where $v_u = 2^{-1/2}v \sin \beta$, $v_{de} = 2^{-1/2}v \cos \beta$, with $v = 2^{-1/4}G_F^{-1/2} = 246$ GeV). A general feature of this ansatz is that it tends to require a rather large top quark mass, m_t . Further recent works have considered more complicated mass matrices [8] and higher-dimension operators [9] which do not, in general, yield (1) (see also Ref. [10]).

In this paper we exhibit a new and quite different model for fermion mass matrices which yields the GJ mass relation (1). We show that this model agrees with experiment, in particular with values of m_t around 135 GeV, in the range suggested by MSSM fits to the LEP data [3, 11]. Our model is defined at the SGUT scale by

$$Y_u = \begin{pmatrix} 0 & A_u & 0 \\ A_u & B_u & 0 \\ 0 & 0 & C_u \end{pmatrix} \quad (5)$$

$$Y_d = \begin{pmatrix} 0 & A_d e^{i\phi} & 0 \\ A_d e^{-i\phi} & B_d & B_d \\ 0 & B_d & C_d \end{pmatrix} \quad (6)$$

$$Y_e = \begin{pmatrix} 0 & A_d & 0 \\ A_d & -3B_d & -3B_d \\ 0 & -3B_d & C_d \end{pmatrix} \quad (7)$$

A natural setting for this ansatz would be an SO(10) SGUT, where one can obtain symmetric (complex) Yukawa matrices by using **10** and **126** Higgs representations to couple to the fermion **16** $\times$ **16** bilinear. If one had a fundamental theory of fermion masses, one would presumably be able to derive the forms of the Y_f and also the values of the parameters from first principles. In the absence of such a theory, we believe that models such as ours can give valuable hints about the underlying physics [12]. Our mass matrix model has seven real parameters A_u , B_u , C_u , A_d , B_d , C_d , and ϕ (plus $\tan \beta$) to describe the nine fermion masses in the u, d, e sectors and the four angles parametrizing the quark mixing matrix V (plus $\tan \beta$);

hence it makes six predictions. (Because of the uncertainties regarding neutrino masses and mixing, we do not consider these here.)

We first show that this ansatz yields the GUT scale GJ mass relation (1). The Y_f , $f = u, d, e$ are diagonalized by unitary transformations U_f : $U_f Y_f U_f^\dagger = \text{diag}\{\lambda_{f,1}, \lambda_{f,2}, \lambda_{f,3}\}$, where $\lambda_{f,j} = \pm m_{f,j}/v_f$ (with the quantities defined at $\mu = m_G$, and $v_d = v_e \equiv v_{de}$). Without loss of generality, one may pick $\lambda_{f,3} > 0$. For Y_d one can take either of the choices $(\lambda_{d,1}, \lambda_{d,2}, \lambda_{d,3}) = (+, -, +), (-, +, +)$; correspondingly, $(\lambda_{e,1}, \lambda_{e,2}, \lambda_{e,3}) = (-, +, +), (+, -, +)$. To leading order in small fermion mass ratios, we find that $C_d = C_e \Rightarrow m_b = m_\tau$. Next, $B_d = \lambda_{d,2} = \mp m_s/v_{de}$ and $\lambda_{e,2} = -3\lambda_{d,2}$, which together imply $m_s = m_\mu/3$. Finally, $A_d = \sqrt{-\lambda_{d,1}\lambda_{d,2}} = \sqrt{m_d m_s}/v_{de}$ and $(Y_e)_{12} = (Y_e)_{21} = \sqrt{-\lambda_{e,1}\lambda_{e,2}} = \sqrt{m_e m_\mu}/v_{de}$ so that the property $(Y_e)_{12} = A_d$ implies $\sqrt{m_e m_\mu} = \sqrt{m_d m_s}$ and hence $m_d = 3m_e$. (These relations hold for either set of sign choices.) This proves our assertion. We will use the choice $(-, +, +)$ for Y_d since it gives a slightly better fit to the data on quark mixing. For Y_u one may also take $(\lambda_{u,1}, \lambda_{u,2}, \lambda_{u,3}) = (+, -, +)$ or $(-, +, +)$, and one finds $C_u = \lambda_{u,3} = m_t/v_u$, $B_u = \lambda_{u,2} = \mp m_c/v_u$, and $A_u = \sqrt{-\lambda_{u,1}\lambda_{u,2}} = \sqrt{m_u m_c}/v_u$. The different sign choices for the $\lambda_{u,j}$ will imply picking ϕ values in the fit which differ by π ; for definiteness, we use $(+, -, +)$ for the $\lambda_{u,j}$.

To test the model, we evolve the Yukawa matrices down from the SGUT scale to the electroweak scale, m_{EW} and diagonalize them there. We shall take $m_{EW} = m_t$, but our results are not very sensitive to this choice. We use the illustrative value $m_t(m_t) = 135$ GeV (i.e., pole mass $M_t = m_t(m_t)[1 + 4\alpha_3(M_t)/(3\pi) + O(\alpha_3(M_t)^2)] \simeq 142$ GeV), consistent with MSSM fits to LEP data [3, 11]. To avoid complications involving further model-dependence, we follow a common simplification [6, 7] of approximating the full SUSY mass spectrum by a single mass scale, m_{SUSY} [13] with $m_{SUSY} = m_t$. Given the theoretical uncertainties associated with the SUSY spectrum and threshold corrections, 1-loop RGE's are sufficiently accurate in the range from m_G to m_{SUSY} . The inputs $\alpha_{em}(m_Z)^{-1} = 127.9 \pm 0.2$ and $\sin^2 \hat{\theta}_W = 0.2325 \pm 0.008$ [14], which give $\alpha_1(m_Z)^{-1} = 58.90 \pm 0.11$ and $\alpha_2(m_Z)^{-1} = 29.74 \pm 0.11$, together with the assumptions that the lightest Higgs has $m_h \simeq m_Z$ and $m_{SUSY} \simeq m_t = 135$ GeV lead to 1-loop unification in the MSSM at $m_G = 1.3 \times 10^{16}$, with $\alpha_G^{-1} = 24.8$. We

take $\alpha_3(m_Z) = 0.125$, consistent with current measurements [3, 15] and with MSSM gauge coupling unification. (Using the 3-loop QCD RGE's [16], this gives $\Lambda_{\overline{MS}}^{(5)} = 302$ MeV and $\alpha_3(m_t) = 0.118$.)

An interesting property of the model which we have established by analytic and numerical solutions of the RGE's is that although the values of the parameters change, the forms of Y_u and Y_e are preserved to high accuracy under evolution from m_G to m_{EW} . For Y_d , the relation $(Y_d)_{21} = (Y_d)_{12}^*$ and the equality $(Y_d)_{22} = (Y_d)_{23}$ are maintained quite accurately, while $|(Y_d)_{32}|$ decreases (by about 35 %) relative to $|(Y_d)_{23}|$. Keeping dominant terms (and with $\dot{f}(t) \equiv (16\pi^2)^{-1}df/dt$, $t = \ln(\mu/m_G)$), the RGE's for the Yukawa matrices thus reduce to RGE's for the nonzero elements, which are

$$\dot{X}_u = X_u(-\sum_j c_{u,j}g_j^2) \quad (8)$$

for $X_u = A_u, B_u$;

$$\dot{C}_u = C_u(-\sum_j c_{u,j}g_j^2 + 6C_u^2) \quad (9)$$

$$\dot{X}_d = X_d(-\sum_j c_{d,j}g_j^2) \quad (10)$$

for $X_d = A_d, (Y_d)_{22}, (Y_d)_{23}$;

$$\dot{Z}_d = Z_d(-\sum_j c_{d,j}g_j^2 + C_u^2) \quad (11)$$

for $Z_d = (Y_d)_{32}, (Y_d)_{33}$; and

$$\dot{X}_e = X_e(-\sum_j c_{e,j}g_j^2) \quad (12)$$

for $X_e = A_e, B_e, C_e$, where for the MSSM in the relevant energy range $\mu > m_{SUSY}$, $(b_1, b_2, b_3) = (\frac{33}{5}, 1, -3)$, $(c_{u,1}, c_{u,2}, c_{u,3}) = (\frac{13}{15}, 3, \frac{16}{3})$, $(c_{d,1}, c_{d,2}, c_{d,3}) = (\frac{7}{15}, 3, \frac{16}{3})$, $(c_{e,1}, c_{e,2}, c_{e,3}) = (\frac{9}{5}, 3, 0)$ [17]. To this order, the running of ϕ is negligibly small.

Manipulating these RGE's, we obtain the solutions

$$\frac{m_b(m_{EW})}{m_\tau(m_{EW})} = \frac{C_d(t_{EW})}{C_e(t_{EW})} = \left(\frac{C_u(t_{EW})}{C_u(t_G)}\right)^{1/6} \rho_C(t_{EW}) \quad (13)$$

where

$$\rho_C(t) = \prod_{j=1}^3 \left(\frac{\alpha_G}{\alpha_j(t)}\right)^{\frac{c_{d,j}-c_{e,j}-(c_{u,j}/6)}{2b_j}} \quad (14)$$

and $t_{EW} = \ln(m_t/m_G)$. We calculate $\rho_C(t_{EW}) = 1.92$. Further, we define $\eta_f = m_f(m_f)/m_f(m_{EW})$ for leptons and heavy quarks, and $\eta_{uds} = m_f(1 \text{ GeV})/m_f(m_{EW})$ for u, d, s and take the value $m_b(m_b) = 4.19 \text{ GeV}$, consistent with Ref. [20]. Using 1-loop QED and 3-loop QCD RGE's, we obtain $\eta_\tau = 1.02$ and $\eta_b = 1.55$, whence $m_b(m_{EW})/m_\tau(m_{EW}) = 1.54$. We successfully fit this ratio by choosing an appropriate value of $C_u(t_G)$. This is a nontrivial test of the ansatz, since it is not, *a priori*, guaranteed that a perturbatively acceptable value of $C_u(t_G)$ would enable one to fit the observed m_b/m_τ mass ratio. We find $C_u(t_{EW})/C_u(t_G) = 0.269$ from (13); substituting this into the solution to the RGE for C_u [18], we get $C_u(t_G) = 4.48$. (This is consistent with the validity of perturbation theory since loop corrections are small: $C_u^2/(16\pi^2) = 0.13$.) We then get $C_u(t_{EW}) = 1.21$ and, from the condition $m_t(m_t) = C_u(t_{EW})v \sin(\beta)/\sqrt{2}$, determine $\tan \beta = 0.84$.

From the RGE's, we obtain the solutions

$$\frac{A_d(t)}{A_e(t)} = \frac{B_d(t)}{B_e(t)} = \rho_{AB}(t) \quad (15)$$

where $(Y_d)_{22}(t) = (Y_d)_{23} \equiv B_d(t)$ and

$$\rho_{AB}(t) = \prod_{j=1}^3 \left(\frac{\alpha_G}{\alpha_j(t)} \right)^{\frac{c_{d,j} - c_{e,j}}{2b_j}} \quad (16)$$

We calculate $\rho_{AB}(t = t_{EW}) = 2.38$. Now

$$\frac{m_s(m_{EW})}{m_\mu(m_{EW})} = \frac{B_d(t_{EW})}{3B_e(t_{EW})} = 0.792 \quad (17)$$

Using $m_\mu(m_{EW}) = 102.8 \text{ MeV}$, the model predicts $m_s(m_{EW}) = 81.4 \text{ MeV}$, i.e., with our 3-loop $\eta_{uds} = 2.94$, $m_s(1 \text{ GeV}) = 239 \text{ MeV}$. This is consistent with the (upper range of the) determination $m_s(1 \text{ GeV}) = 175 \pm 55 \text{ MeV}$ [20]. Further we find

$$\frac{(Y_d)_{32}(t)}{(Y_d)_{23}(t)} = \left(\frac{C_u(t)}{C_u(t_G)} \right)^{\frac{1}{6}} \prod_{j=1}^3 \left(\frac{\alpha_G}{\alpha_j(t)} \right)^{-\frac{c_{u,j}}{12b_j}} \quad (18)$$

We use the results $A_d(t)v_{de} = \sqrt{m_d(t)m_s(t)}$, $A_e(t)v_{de} = \sqrt{m_e(t)m_\mu(t)}$ and eq. (15) to find $m_d(m_{EW})/m_e(m_{EW}) = 7.13$. Using $m_e(m_{EW}) = 0.4876 \text{ MeV}$, this yields $m_d(m_{EW}) = 3.48 \text{ MeV}$, and hence $m_d(1 \text{ GeV}) = 10.2 \text{ MeV}$, in agreement with the value $m_d(1 \text{ GeV}) = 8.9 \pm 2.6 \text{ MeV}$ [20].

Given that Y_u retains its form to high accuracy under the evolution from m_G to m_{EW} , the two remaining parameters in Y_u are determined via the relations $B_u(t_{ew})v_u = -m_c(t)$ and $A_u(t_{ew})v_u = \sqrt{m_u(t)m_c(t)}$. From $m_c(m_c) = 1.27 \pm 0.05$ GeV [20] we calculate $\eta_c = 2.32$, whence $m_c(m_{EW}) = 0.547 \pm 0.022$ GeV. With $m_u(1 \text{ GeV}) = 5.1 \pm 1.5$ MeV we get $m_u(m_{EW}) = 1.7 \pm 0.5$ MeV. Diagonalizing at m_{EW} , we have $v_u U_{u,L} Y_u U_{u,L}^\dagger = M_u$, $v_d^2 U_{d,L} Y_d Y_d^\dagger U_{d,R}^\dagger = M_d^2$, whence $V = U_{u,L} U_{d,L}^\dagger$ for the Cabibbo-Kobayashi-Maskawa (CKM) matrix V [21]. To leading order $|V_{us}| \simeq |\sqrt{m_u/m_c} - e^{i\phi} \sqrt{m_d/m_s}|$ (with quantities evaluated at $\mu = m_{EW}$ [21]). We determine $\phi = 75.5^\circ$ by fitting $|V_{us}|$. Quoting as usual the rephasing-invariant $|V_{jk}|$ values, we get

$$\{|V_{ij}|\} = \begin{pmatrix} 0.9753 & 0.221 & 0.002 \\ 0.221 & 0.9748 & 0.030 \\ 0.006 & 0.030 & 0.9995 \end{pmatrix} \quad (19)$$

in agreement with current data [14, 19] (with $|V_{cb}|$ in the lower range of acceptable values). For the reparametrization-invariant CP violation parameter J [22], we get $J = 0.95 \times 10^{-5}$, again in agreement with data. Further details will be given elsewhere [23].

References

- [1] Reviews include H. Nilles, Phys. Repts. **110**, 1 (1984); H. Haber and G. Kane, Phys. Repts. **117** (1984); for recent discussions of gauge coupling unification and mass limits, and references to the literature, see, e.g., *Ten Years of SUSY Confronting Experiment*, CERN-TH.6707/92-PPE/92-180 and *Proceedings of SUSY-93*, Northeastern University.
- [2] A recent review is T. Appelquist, in *Proc. XXVI Int. Conf. on High Energy Physics*, Dallas, 1992 (A.I.P., 1993), p. 1333.
- [3] L. Rolandi, in *Proc. XXVI Conf. H. E. Physics*, 1992 *op. cit.*, p. 56.
- [4] H. Georgi and C. Jarlskog (GJ), Phys. Lett. **B86**, 297 (1979).
- [5] H. Arason, D. Castaño, B. Keszthelyi, S. Mikaelian, E. Piard, P. Ramond, and B. Wright Phys. Rev. Lett. **67**, 2933 (1991); Phys. Rev. D **46**, 3945 (1992); H. Arason, D. Castaño, P. Ramond, and E. Piard, Phys. Rev. D **47**, 232 (1993).

- [6] S. Dimopoulos, L. Hall, and S. Raby, Phys. Rev. Lett.**68**, 1984 (1992); Phys. Rev. D**45**, 4195 (1992); Anderson, S. Raby, S. Dimopoulos, L. Hall, Phys. Rev. D**47**, R3702 (1993).
- [7] V. Barger, M. Berger, T. Han, and M. Zralek, Phys. Rev. Lett.**68**, 3394 (1992); V. Barger, M. Berger, and P. Ohmann, Phys. Rev. D**47**, 1093 (1992).
- [8] G. Giudice, Mod. Phys. Lett. **A7**, 2429 (1992).
- [9] L. Hall, LBL-33677 (hep-ph/9303217); S. Raby, OHSTPY-HEP-T-93007 (hep-ph/9306335).
- [10] P. Ramond, R. Roberts, and G. G. Ross, preprint RAL-93-010/UFIFT-93-010 (hep-ph/9303320) considers SGUT quark mass matrices.
- [11] G. Altarelli, R. Barbieri, and S. Jadach, Nucl. Phys. B**369**, 3 (1992); erratum **B376**, 444 (1992); R. Barbieri, M. Frigeni, and F. Caravaglios, Phys. Lett. B**279**, 169 (1992).
- [12] It has often been speculated that a given ansatz may arise in a technically natural way due to global symmetries. However, global symmetries are likely to be broken by quantum gravity (see, e.g., the review by M. Dine, in *Proc. XXVI Conf. H. E. Physics*, 1992 *op. cit.*, p. 1386). Discrete symmetries do arise naturally in string theory, where they are typically local. Perhaps the zeroes in mass matrices should really be regarded as only good approximations to nonzero quantities dynamically determined by a more fundamental theory (e.g. R. Shrock, Phys. Rev. D**45** R10 (1992)). Our model specifically assumes that the underlying theory yields (i) zeroes as indicated; (ii) the relation $|(Y_d)_{22}| = |(Y_d)_{23}| = |(Y_d)_{32}|$ and (iii) $Im((Y_d)_{jk}) = 0$, $jk = 23, 32$. We do not assume that the properties which lead to (5)-(7) hold below the SGUT level. Interestingly, the zeroes, the reality of $(Y_d)_{jk}$, $jk = 23, 32$, and the relation $(Y_d)_{22} = (Y_d)_{23}$ still hold to very good accuracy down to t_{EW} . We thank L. Lavoura for a helpful comment.
- [13] To do significantly better than the assumption of a single m_{SUSY} would require a self-consistent evolution of the full theory down from the SGUT level, which would predict

both the supersymmetric particle mass spectrum and the usual Yukawa mass matrices, while addressing issues such as the “ μ problem”, doublet-triplet splitting, flavor-changing neutral operators, etc; inevitably, this would involve more parameters and model-dependence. The effect of using $m_{SUSY} = 1$ TeV as well as $m_{SUSY} = m_t = 150$ GeV has been studied in Ref. [7].

- [14] Particle Data Group, *Review of Particle Properties*, Phys. Rev. D**45** III.65 (1992).
- [15] S. Bethke, in *Proc. XXVI Int. Conf. on High Energy Physics*, Dallas, 1992 (A.I.P., 1993), p. 81; R. K. Ellis, in *DPF 92*, C. Albright, ed. (World Scientific, 1993), p. 167.
- [16] O. Tarasov, A. Vladimirov, and A. Zharkov, Phys. Lett. B**93**, 429 (1980); S. Gorishnii, A. Kataev, and S. Larin, Yad. Fiz. **40**, 517 (1984) [Sov. J. Nucl. Phys. **40**, 329 (1984)].
- [17] K. Inoue, A. Kakuto, H. Komatsu, and S. Takeshita, Prog. Theor. Phys. **67**, 1889 (1992) and references therein.
- [18] Since $C_u \gg$ all other terms in the Y_f , the dominant terms in the 1-loop RGE for C_u (9) are the same for our model and that of (2)-(4); the solution is $(C_u(t)/C_u(t_G))^2 = \rho(t)/[1 + 3(2\pi)^{-2}C_u^2(t_G)I(t)]$, where $\rho(t) = \prod_{j=1}^3(\alpha_G/\alpha_j(t))^{c_{u,j}/b_j}$ and $I(t) = \int_t^{t_G} \rho(t')dt'$. For our values of $\alpha_3(m_Z)$ and m_{EW} , we get $\rho(t_{EW}) = 13.2$ and $I(t_{EW}) = 119$.
- [19] F. Muheim, in *Proc. DPF 92, op. cit.*, v. 1, p. 614; A. Bean et al. (CLEO collab.) Phys. Rev. Lett.**70**, 2681 (1993). CLEO II finds $|V_{ub}/V_{cb}| = 0.085 \pm 0.035$, which, with $|V_{cb}| = 0.043 \pm 0.011$, gives $|V_{ub}| \simeq 0.0037 \pm 0.0018$.
- [20] J. Gasser and H. Leutwyler, Phys. Repts. **87**, 77 (1982).
- [21] The running of V from m_{EW} to the low energies where its elements have been measured can be neglected.
- [22] C. Jarlskog, Phys. Rev. Lett.**55**, 1039 (1985); Z. f. Physik **29**, 491 (1985).
- [23] A. Kusenko and R. Shrock, to appear.